

## CREB1

**Reactivity:** Human Mouse Rat Monkey

**Tested applications:** WB IHC IF IP FC

**Recommended Dilution:** WB 1:1000 - 1:4000 IHC 1:50 - 1:200 IF 1:50 - 1:100 IP 1:20 - 1:50  
FC 1:20 - 1:50

**Calculated MW:** 35kd

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human CREB1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

hikq

**Synonym:**

CREB ; CREB1; CREB 1

**Catalog #:** A0011

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 1385

**Isotype:** IgG

**Swiss Prot:** P16220

**Purity:** Affinity purification

For research use only.

**Background:**

CREB1 is a bZIP transcription factor that activates target genes through cAMP response elements. CREB1 is able to mediate signals from numerous physiological stimuli, resulting in regulation of a broad array of cellular responses. While CREB1 is expressed in numerous tissues, it plays a large regulatory role in the nervous system. CREB1 is believed to play a key role in promoting neuronal survival, precursor proliferation, neurite outgrowth, and neuronal differentiation in certain neuronal populations (1-3). Additionally, CREB1 signaling is involved in learning and memory in several organisms (4-6). CREB1 is able to selectively activate numerous downstream genes through interactions with different dimerization partners. CREB1 is activated by phosphorylation at Ser133 by various signaling pathways including Erk, Ca2+, and stress signaling. Some of the kinases involved in phosphorylating CREB1 at Ser133 are p90RSK, MSK, CaMKIV, and MAPKAPK-2 (7-9).

**To place an order, please [Click HERE](#).**