

TSC1

Reactivity: Human Mouse

Tested applications: WB

Recommended Dilution: WB 1:500 - 1:1000

Calculated MW: 130kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human TSC1

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b

Synonym:

KIAA0243; LAM; MGC86987; TSC

Catalog #: A0721

Antibody Type:

Monoclonal Antibody

Species: Mouse

Gene ID: 7248

Isotype: IgG

Swiss Prot: Q92574

Purity: Affinity purification

For research use only.

Background:

TSC1, also named as KIAA0243 and TSC, is implicated as a tumor suppressor. It is involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling. In complex with TSC2, TSC1 inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. It seems not to be required for TSC2 GAP activity towards RHEB. It is involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling. Defects in TSC1 are the cause of tuberous sclerosis complex (TSC). Defects in TSC1 may be a cause of focal cortical dysplasia of Taylor balloon cell type (FCDBC). The antibody is specific to TSC1.

To place an order, please [Click HERE](#).