

HDAC5

Reactivity: Human Mouse

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:50

Calculated MW: 122kDa

Observed MW: Refer to figures

Immunogen:

A synthetic peptide of human HDAC5

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fp

Synonym:

HD5; NY-CO-9;

Catalog #: A7189

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 10014

Isotype: IgG

Swiss Prot: Q9UQL6

Purity: Affinity purification

For research use only.

Background:

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).