www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## KPNA6

Reactivity: Human Mouse

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:60kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human KPNA6

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

IPOA7; KPNA7;

Catalog #:A7363

Gene ID:23633

Isotype:IgG

Swiss Prot: O60684 Purity: Affinity purification

For research use only.

## Background:

Nucleocytoplasmic transport, a signal- and energy-dependent process, takes place through nuclear pore complexes embedded in the nuclear envelope. The import of proteins containing a nuclear localization signal (NLS) requires the NLS import receptor, a heterodimer of importin alpha and beta subunits also known as karyopherins. Importin alpha binds the NLS-containing cargo in the cytoplasm and importin beta docks the complex at the cytoplasmic side of the nuclear pore complex. In the presence of nucleoside triphosphates and the small GTP binding protein Ran, the complex moves into the nuclear pore complex and the importin subunits dissociate. Importin alpha enters the nucleoplasm with its passenger protein and importin beta remains at the pore. The protein encoded by this gene is a member of the importin alpha family.

To place an order, please Click HERE.







