www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

RARA

Reactivity: Human

Tested applications: WB IP ChIP

Recommended Dilution:WB 1:500 - 1:2000 IP 1:20 - 1:50 ChIP 1:20 - 1:50

Calculated MW:51kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human RARA

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

kt

Synonym:

RAR; NR1B1;

Background:

This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoeisis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus.

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:5914

Isotype:IgG

Swiss Prot:P10276

Purity: Affinity purification

For research use only.







