

## BAX

**Reactivity:** Human

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:2000

**Calculated MW:** 21kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human BAX

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bhi

**Synonym:**

BCL2L4;

**Catalog #:** A7626

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 581

**Isotype:** IgG

**Swiss Prot:** Q07812

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene.

*To place an order, please [Click HERE](#).*