

HLA-DRB1

Reactivity: Human

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW: 30kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human HLA-DRB1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

SS1; DRB1; DRw10; HLA-DRB; HLA-DR1B;

Catalog #: A7685

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 3123

Isotype: IgG

Swiss Prot: P04229

Purity: Affinity purification

For research use only.

Background:

HLA-DRB1 belongs to the HLA class II beta chain paralogs. The class II molecule is a heterodimer consisting of an alpha (DRA) and a beta chain (DRB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins.

Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa. It is encoded by 6 exons. Exon one encodes the leader peptide; exons 2 and 3 encode the two extracellular domains; exon 4 encodes the transmembrane domain; and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Hundreds of DRB1 alleles have been described and typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogs DRB3, DRB4 and DRB5. DRB1 is present in all individuals. Allelic variants of DRB1 are linked with either none or one of the genes DRB3, DRB4 and DRB5. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9.

To place an order, please [Click HERE](#).