## MOCS<sub>1</sub>

Reactivity: Human Mouse

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:70kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MOCS1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

MIG11; MOCOD; MOCODA;

**Antibody Type:** 

Polyclonal Antibody

Species:Rabbit

Gene ID:4337

Isotype:IgG

Swiss Prot:Q9NZB8

Purity: Affinity purification

For research use only.

## Background:

Molybdenum cofactor biosynthesis is a conserved pathway leading to the biological activation of molybdenum. The protein encoded by this gene is involved in this pathway. This gene was originally thought to produce a bicistronic mRNA with the potential to produce two proteins (MOCS1A and MOCS1B) from adjacent open reading frames. However, only the first open reading frame (MOCS1A) has been found to encode a protein from the putative bicistronic mRNA, whereas additional splice variants, whose full-length natures have yet to be determined, are likely to produce a fusion between the two open reading frames. This gene is defective in patients with molybdenum cofactor deficiency, type A. A related pseudogene has been identified on chromosome 16.

To place an order, please Click HERE.





