

PTTG1

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW: 22kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human PTTG1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

EAP1; PTTG; HPTTG; TUTR1;

Catalog #: A8307

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 9232

Isotype: IgG

Swiss Prot: O95997

Purity: Affinity purification

For research use only.

Background:

The encoded protein is a homolog of yeast securin proteins, which prevent separins from promoting sister chromatid separation. It is an anaphase-promoting complex (APC) substrate that associates with a separin until activation of the APC. The gene product has transforming activity in vitro and tumorigenic activity in vivo, and the gene is highly expressed in various tumors. The gene product contains 2 PXXP motifs, which are required for its transforming and tumorigenic activities, as well as for its stimulation of basic fibroblast growth factor expression. It also contains a destruction box (D box) that is required for its degradation by the APC. The acidic C-terminal region of the encoded protein can act as a transactivation domain. The gene product is mainly a cytosolic protein, although it partially localizes in the nucleus. Three transcript variants encoding the same protein have been found for this gene.

To place an order, please [Click HERE](#).