www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IL5RA

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:48kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human IL5RA

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

IL5R; CD125; CDw125; HSIL5R3

Background:

The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported.

To place an order, please Click HERE.

Polyclonal Antibody Species: Rabbit Gene ID:3568 Isotype:IgG Swiss Prot:Q01344 Purity: Affinity purification

For research use only.





