

## Phospho-NFKB2-S870

**Reactivity:** Human

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:2000

**Calculated MW:** 120kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A phospho specific peptide corresponding to residues surrounding S870 of human NFKB2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

kop

**Synonym:**

p52; p105; H2TF1; LYT10; CVID10; LYT-10; NF-kB2;

**Catalog #:** AP0419

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 4791

**Isotype:** IgG

**Swiss Prot:** Q00653

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.

**To place an order, please [Click HERE](#).**