

## MIP-1 Gamma Mouse

**Description:** MIP-1 gamma Mouse Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 101 amino acids and having a molecular mass of 11.6 kDa. The MIP-1 gamma is purified by proprietary chromatographic techniques.

**Catalog #:** CHPS-264

For research use only.

**Synonyms:** CCL9/10, MRP2, CCF18.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** QITHATETKE VQSSLKAQQG LEIEMFHMGF QDSSDCCLSY  
NSRIQCSRFI GYFPTSGGCT RPIIFISKR GFQVCANPSD RRVQRCIERL EQNSQPRTYK Q.

**Purity:** Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

The MIP-1 gamma was lyophilized from 1xPBS solution pH-7.4.

**Stability:**

Lyophilized MIP-1 gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL9/10 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized MIP-1 gamma in sterile 18M-cm H<sub>2</sub>O not less than 100

**Introduction:**

Mouse MIP-1 gamma is 75% identical in its amino acid composition as compared to the rat species. MIP-1 gamma is a CC chemokine localized in murine blood and a widespread range of murine tissues, without having an identified human homolog. MIP-1 gamma signals through the CCR1 receptor. MIP-1 gamma chemoattracts neutrophils and also inhibits colony formation of bone marrow myeloid immature progenitors. MIP-1 gamma has six cysteines including the four highly conserved cysteine residues present in CC chemokines.

**Biological Activity:**

Defined by its ability to chemoattract human neutrophils using a concentration range of 0.1-10 ng/ml, corresponding to a Specific Activity of 10,000-100,000IU/mg.

**To place an order, please [Click HERE](#).**