www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Acrp30 Mouse, HEK

Description: The Acrp30 Mouse Recombinant is fused with FLAG tag having a total Mw of 26kDa.

Catalog #:CYPS-442

Synonyms: Acrp30, AdipoQ, GBP-28, APM-1, ACDC.

For research use only.

Source: HEK293 (Human embryonic kidney cell line).

Amino Acid Sequence: EDDVTTTEEL APALVPPPKG TCAGWMAGIP GHPGHNGTPG RDGRDGTPGE KGEKGDAGLL GPKGETGDVG MTGAEGPRGF PGTPGRKGEP GEAAYMYRSA FSVGLETRVT VPNVPIRFTK IFYNQQNHYD GSTGKFYCNI PGLYYFSYHI TVYMKDVKVS LFKKDKAVLF TYDQYQEKNV DQASGSVLLH LEVGDQVWLQ VYGDGDHNGL YADNVNDSTF TGFLLYHDTN DY

Purity: Purity of adiponectin mouse recombinant is >98% (HPLC and SDS PAGE analyzed).

Formulation:

Sterile filtered and lyophilized from 0.5 mg/ml in PBS buffer.

Stability:

Store lyophilized APM-1 at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely.

Introduction:

Adiponectin is an important negative regulator in hematopoiesis and immune systems. It may be involved in ending inflammatory responses through its inhibitory functions. Inhibits endothelial nf-kappa-b signaling through a camp-dependent pathway. Inhibits tnf-alpha- induced expression of endothelial adhesion molecules. Involved in the control of fat metabolism and insulin sensitivity.

Biological Activity:

Full-length mouse adiponectin activates AMP-activated protein kinase in hepatocyte and activates AMPK in HepG2 human hepatocytes at a concentration of 1000ng/ml corresponding to a Specific Activity of 1000IU/mg. Mouse adiponectin mammalian cell derived inhibits glucose production as shown by in-vitro gluconeogenesis assay in primary rat hepatocytes.

To place an order, please Click HERE.





