

## DHFR Mouse

**Description:** DHFR Mouse Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 207 amino acids (1-187 a.a.) and having a molecular mass of 23.8kDa. The DHFR is purified by proprietary chromatographic techniques.

Catalog #:ENPS-076

For research use only.

**Synonyms:** Dihydrofolate reductase, DHFR, DHFRP1, AA607882, AI662710, AW555094, 8430436I03Rik.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered colorless solution.

**Amino Acid Sequence:** MGSSHHHHHH SSGLVPRGSH MVRPLNCIVA VSQNMGIGKN  
GDLWPPLRN EFKYFQRTT TSSVEGQNL VIMGRKTWFS IPEKNRPLKD RINIVLSREL  
KEPPRGAFHL AKSLDDALRL IEQPELASKV DMVWIVGGSS VYQEAMNQPG HLRLFVTRIM  
QEFESDTFFP EIDLGKYKLL PEYPGVLSEV QEEKGIKYKF EYVEKKD.

**Purity:** Greater than 95.0% as determined by SDS-PAGE.

### Formulation:

The DHFR solution (1 mg/ml) contains 20mM Tris-HCl buffer(pH 8.0), 10% glycerol, 2mM DTT and 0.1M NaCl.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

Dihydrofolate reductase (DHFR) is an enzyme that reduces dihydrofolic acid to tetrahydrofolic acid, with NADPH as electron donor that can be converted to the kinds of tetrahydrofolate cofactors applied in 1-carbon transfer chemistry. DHFR converts dihydrofolate into tetrahydrofolate, which is a methyl group shuttle required for the de novo synthesis of purines, thymidyl acid, and specific amino acids. Even though the functional DHFR gene is mapped to chromosome 5, numerous intronless processed pseudogenes or dihydrofolate reductase-like genes are identified on separate chromosomes. DHFR deficiency is associated with megaloblastic anemia. DHFR knockdown plays a role in the anticancer activity of 2-hydroxyoleic acid. DHFR gene insertion/deletion polymorphism is linked to variation in serum and red blood cell folate concentrations in women.

### Biological Activity:

Specific activity is >0.2 units/mg, in which one unit will convert 1.0 umole of 7,8-dihydrofolate and beta-NADPH to 5,6,7,8-tetrahydrofolate and beta-NADP per min at pH 6.5 at 25°C.

**To place an order, please [Click HERE](#).**