

PPA E.Coli

Description: PPA E.Coli Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 196 amino acids (1-176 a.a.) and having a molecular mass of 21.9kDa. PPA is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: ENPS-156

For research use only.

Synonyms: Inorganic pyrophosphatase, Pyrophosphate phospho-hydrolase, PPase, ppa, b4226, JW4185.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MSLLNVPAGK DLPEDIYVVI
EIPANADPIK YEIDKESGAL FVDRFMSTAM FYPCNYGYIN HTLSLDGDPV DVLVPTYPL
QPGSVIRCRP VGLKMTDEA GEDAKLVAVP HSKLSKEYDH IKDVNDLPEL LKAQIAHFFE
HYKDLEKGGW VKVEGWENAE AAKAEIVASF ERAKNK.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

PPA protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 1mM DTT, 10% glycerol and 50mM NaCl.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Inorganic pyrophosphatase (ppa) is a member of the Ppase family. PPA is an enzyme which catalyzes the conversion of one molecule of pyrophosphate to two phosphate ions. Since this is a highly exergonic reaction, it can therefore be coupled to unfavorable biochemical transformations in order to drive these transformations to completion. The role of the PPA enzyme is a critical one in the lipid metabolism (including lipid synthesis and degradation), calcium absorption and bone formation, DNA synthesis, as well as other biochemical transformations.

To place an order, please [Click HERE](#).