

MUTM E.Coli

Description: MUTM Recombinant produced in E. coli is a single polypeptide chain containing 289 amino acids (1-269) and having a molecular mass of 32.4kDa. MUTM is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: ENPS-596

For research use only.

Synonyms: Formamidopyrimidine-DNA glycosylase, FPG.

Source: E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MPELPEVETS RRGIEPHLVG
ATILHAVVRN GRLRWPVSEE IYRLSDQPV LSVQRRAKYLL LELPEGWIII HLGMSGSLRI
LPEELPPEKH DHVDLMSNG KVLRYTDPRR FGAWLWTKEL EGHNVLTHLG PEPLSDDFNG
EYLHQKCAKK KTAIKPWLM D NKLVVGVGNI YASESLFAAG IHPDRLASSL SLAECELLAR
VIKAVLLRSI EQ

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

The MUTM solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 100mM NaCl, 1mM DTT and 20% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

MUTM is a base excision repair enzyme that identifies and eliminates a large variety of oxidized purines from correspondingly impaired DNA. MUTM is nondismissable and essential to remove quickly its substrate lesions on the chromosome. MUTM, additionally, mends a large number of the lesions recognized by Endo III, signifying that MUTM takes a prominent part in the overall repair of both purine damage and pyrimidine damage in vivo.

To place an order, please [Click HERE](#).