

MSH2

Reactivity:Human

Tested applications:WB IF FC

Recommended Dilution:WB 1:500 - 1:1000 IF 1:20 - 1:50 FC 1:20 - 1:50

Calculated MW:105kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human MSH2

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

be

Synonym:

MSH2;COCA1;FCC1;HNPCC;HNPCC1;LCFS2

Catalog #:A0064

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:4436

Isotype:IgG

Swiss Prot:P43246

Purity:Affinity purification

For research use only.

Background:

The DNA mismatch repair system (MMR) repairs post-replication DNA, inhibits recombination between non-identical DNA sequences and induces both checkpoint and apoptotic responses following certain types of DNA damage (1). MSH2 (MutS homologue 2) forms the hMutS- dimer with MSH6 and is an essential component of the mismatch repair process. hMutS- is part of the BRCA1-associated surveillance complex (BASC), a complex that also contains BRCA1, MLH1, ATM, BLM, PMS2 proteins and the Rad50-Mre11-NBS1 complex (2). Mutations in MSH2 have been found in a large proportion of hereditary non-polyposis colorectal cancer (Lynch Syndrome), the most common form of inherited colorectal cancer in the Western world (3). Mutations have also been associated with other sporadic tumors.

To place an order, please [Click HERE](#).