

SPHK1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:43kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human SPHK1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

ab

Synonym:

PHK1; SPHK

Catalog #:A0139

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8877

Isotype:IgG

Swiss Prot:Q9NYA1

Purity:Affinity purification

For research use only.

Background:

Sphingosine kinases (SPHKs) catalyze the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. Together with other sphingolipid metabolizing enzymes, SPHKs regulate the balance of the lipid mediators, ceramide, sphingosine, and S1P (1-4). Two distinct SPHK isoforms, SPHK1 and SPHK2, have been cloned and characterized (5,6). SPHK1 and SPHK2 are highly conserved and diversely expressed (7,8). The SPHKs are activated by G-protein-coupled receptors, receptor tyrosine kinases, immunoglobulin receptors, cytokines, and other stimuli (9-12). The molecular mechanisms by which SPHK1 and SPHK2 are specifically regulated are complex and only partially understood.

To place an order, please [Click HERE](#).