

VASP

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF IP FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 IP 1:20 - 1:50
FC 1:20 - 1:50

Calculated MW:40kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human VASP

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

Vasodilator-stimulated phosphoprotein;VASP;

Catalog #:A0166

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7408

Isotype:IgG

Swiss Prot:P50552

Purity:Affinity purification

For research use only.

Background:

Vasodilator-stimulated phosphoprotein (VASP) was originally characterized as a substrate of both cGMP- and cAMP-dependent kinases (PKG and PKA, or cGPK and cAPK, respectively) (1). It is now believed that VASP belongs to the Ena/VASP family of adaptor proteins linking the cytoskeletal system to the signal transduction pathways and that it functions in cytoskeletal organization, fibroblast migration, platelet activation and axon guidance (2,3). Three phosphorylation sites, Ser157, Ser239, and Thr278, have been identified. Ser239 is the major PKG phosphorylation site while Ser157 is the major PKA phosphorylation site (4). Evidence suggests that VASP phosphorylation reduces its association with actin and has a negative effect on actin polymerization (5). Phosphorylation at Ser239 of VASP is a useful marker for monitoring PKG activation and signaling (6,7).

To place an order, please [Click HERE](#).