

## BCL2L1

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:** 26kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human BCL2L1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bdi

**Synonym:**

BCL2L1;BCL-XL/S;BCL2L;BCLX;Bcl-X;DKFZp781P2092;bcl-xL;bcl-xS;BCLXS

**Catalog #:** A0209

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 598

**Isotype:** IgG

**Swiss Prot:** Q07817

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Two alternatively spliced transcript variants, which encode distinct isoforms, have been reported. The longer isoform acts as an apoptotic inhibitor and the shorter form acts as an apoptotic activator. [provided by RefSeq]

**To place an order, please [Click HERE](#).**