

## CRP

---

**Reactivity:**Human Mouse

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**25kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human CRP

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CRP;MGC149895;MGC88244;PTX1 ;

**Catalog #:**A0224

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**1401

**Isotype:**IgG

**Swiss Prot:**P02741

**Purity:**Affinity purification

For research use only.

**Background:**

Human c-Reactive protein (CRP) was discovered in serum of patients as a nonantibody precipitin of the C-polysaccharide from pneumococci(1). During acute infection, the amounts of circulating CPR increases 100-1000 fold (2). Serum CRP values are routinely measured to detect or monitor many human diseases and minor elevations of C-reactive protein are predictive of cardiovascular events in patients with coronary heart disease (3). CRP promotes host defense, is an efficient activator of the classical C cascade, and shown to be opsonic (4).The opsonic and C activation activities are responsible for the nonspecific host defense activities providing protection in the preimmune stage of acute infections. Additionally, IL-1 and IL-6 are known to stimulate CRP biosynthesis (5).

**To place an order, please [Click HERE](#).**