

## HSPB1

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IF

**Recommended Dilution:**WB 1:500 - 1:2000 IF 1:20 - 1:100

**Calculated MW:**27kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human HSPB1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

HSPB1;CMT2F;DKFZp586P1322;HMN2B;HS.76067;HSP27;HSP28;Hsp25;SRP27 ;

**Catalog #:**A0240

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**3315

**Isotype:**IgG

**Swiss Prot:**P04792

**Purity:**Affinity purification

For research use only.

**Background:**

Heat shock protein (HSP) 27 is one of the small HSPs that are constitutively expressed at different levels in various cell types and tissues. Like other small heat shock proteins, HSPB1 is regulated at both the transcriptional and posttranslational levels (1). In response to stress, the expression level of HSPB1 increases several-fold to confer cellular resistance to the adverse environmental change. HSPB1 is phosphorylated at Ser15, Ser78 and Ser82 by MAPKAP kinase 2 as a result of the activation of the p38 MAP kinase pathway (2,3). Phosphorylation of HSPB1 causes a change in its tertiary structure, which shifts from large homotypic multimers to dimers and monomers (4). It has been shown that phosphorylation and increased concentration of HSPB1 modulates actin polymerization and reorganization (5,6).

**To place an order, please [Click HERE](#).**