

NAMPT

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:52kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human NAMPT

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

NAMPT;1110035O14Rik;DKFZp666B131;MGC117256;PBEF;PBEF1;VF;VISFATIN ;

Catalog #:A0256

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:10135

Isotype:IgG

Swiss Prot:P43490

Purity:Affinity purification

For research use only.

Background:

Nicotinamide phosphoribosyltransferase(NAMPT) has two usual synonyms termed Visfatin and PBEF. Its primary role is to catalyze the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, an intermediate in the biosynthesis of NAD, which is the rate limiting component in the mammalian NAD biosynthesis pathway. NAMPT is localized in cytoplasm and expressed in large amounts in bone marrow, liver tissue, and muscle tissues. NAMPT inhibits neutrophil apoptosis in experimental inflammation and clinical sepsis. NAMPT levels are altered in plasma of patients with type 2 diabetes mellitus (T2DM), and it is now evidenced that NAMPT may plays a role in lipid metabolism.

To place an order, please [Click HERE](#).