

ABL1

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW:123kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human ABL1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

br

Synonym:

ABL1;ABL;JTK7;bcr/abl;c-ABL;p150;v-abl ;

Catalog #:A0282

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:25

Isotype:IgG

Swiss Prot:P00519

Purity:Affinity purification

For research use only.

Background:

The c-Abl proto-oncogene encodes a nonreceptor protein tyrosine kinase that is ubiquitously expressed and highly conserved in metazoan evolution. c-Abl protein is distributed in both the nucleus and the cytoplasm of cells. It is implicated in regulating cell proliferation, differentiation, apoptosis, cell adhesion and stress responses (1-3). c-Abl kinase activity is increased in vivo by diverse physiological stimuli including integrin activation, PDGF stimulation and binding to c-Jun, Nck and RFX1 (2,4). The in vivo mechanism of regulation of c-Abl kinase activity is not completely understood. Tyr245 is located in the linker region between the SH2 and catalytic domains. This positioning is conserved among Abl family members. Phosphorylation of Tyr245 is involved in the activation of c-Abl kinase (5). In addition, phosphorylation of Tyr412, which is located in the kinase activation loop of c-Abl, is required for kinase activity (6).

To place an order, please [Click HERE](#).