

HSPA1A

Reactivity:Human Mouse

Tested applications:WB IF

Recommended Dilution:WB 1:500 - 1:2000 IF 1:50 - 1:200

Calculated MW:70kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human HSPA1A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bhi

Synonym:

HSP72; HSPA1; HSP70I; HSP70-1; HSP70-1A; HSPA1A

Catalog #:A0284

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3303

Isotype:IgG

Swiss Prot:P08107

Purity:Affinity purification

For research use only.

Background:

HSPA1A and HSP90 are molecular chaperones expressed constitutively under normal conditions to maintain protein homeostasis and are induced upon environmental stress (1). Both HSPA1A and HSP90 are able to interact with unfolded proteins to prevent irreversible aggregation and catalyze the refolding of their substrates in an ATP- and co-chaperone-dependent manner (1). HSPA1A has a broad range of substrates including newly synthesized and denatured proteins, while HSP90 tends to have a more limited subset of substrates, most of which are signaling molecules. HSPA1A and HSP90 often function collaboratively in a multi-chaperone system, which requires a minimal set of co-chaperones: HSP40, Hop, and p23 (2,3). The co-chaperones either regulate the intrinsic ATPase activity of the chaperones or recruit chaperones to specific substrates or subcellular compartments (1,4). When the ubiquitin ligase CHIP associates with the HSPA1A/HSP90 complex as a cofactor, the unfolded substrates are subjected to degradation by the proteasome (4). The biological functions of HSPA1A/HSP90 extend beyond their chaperone activity. They are essential for the maturation and inactivation of nuclear hormones and other signaling molecules (1,3). They also play a role in vesicle formation and protein trafficking (2).

To place an order, please [Click HERE](#).