

CD40LG

Reactivity:Human

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:29kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human CD40LG

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bp

Synonym:

CD40LG;CD154;CD40L;HIGM1;IGM;IMD3;T-BAM;TNFSF5;TRAP;gp39;hCD40L ;

Catalog #:A0327

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:959

Isotype:IgG

Swiss Prot:P29965

Purity:Affinity purification

For research use only.

Background:

CD40 Ligand (CD40L), a member of the TNF superfamily of ligands, is expressed as either a membrane-bound or soluble homotrimer (1). Both membrane-bound and soluble forms of CD40L have biological activity (1,2). CD40L is expressed primarily on activated T cells; however, mast cells, basophils, and NK cells may also express CD40L (1). CD40L functions as an important T cell co-stimulatory molecule and enhances T-dependent B cell responses (1,2). CD40L binds CD40, which is a receptor expressed on B cells, dendritic cells, and macrophages (1). Binding of CD40L to CD40 leads to the recruitment of TRAF proteins (TRAFs 2,3,5, and 6) resulting in the activation of the JNK and NF-B pathways (1,2).

To place an order, please [Click HERE](#).