

TGFA

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW:17kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human TGFA

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

abp

Synonym:

TFGA; Pro-TGF-alpha; TGF-alpha

Catalog #:A0337

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7039

Isotype:IgG

Swiss Prot:P01135

Purity:Affinity purification

For research use only.

Background:

Transforming growth factor alpha (TGF-alpha) is a member of the epidermal growth factor (EGF) family, sharing the same receptor, EGFR, and regulating cell proliferation, survival, and differentiation (1). Members of the family share an EGF-like domain of 45-60 amino acids characterized by the conservation of six regularly spaced cysteins, forming three disulfide bonds that function as their receptor binding domain. TGF-alpha was initially discovered in the media of retrovirally transformed fibroblasts, and its name comes from its ability to induce transformation in cultured fibroblasts (2). This transforming activity was later shown to require TGF-beta, which potentiates the activity of TGF-alpha through a separate receptor (3). Soluble TGF-alpha is released from its membrane-bound precursor, pro-TGF-alpha, following proteolytic cleavage, but the membrane bound precursor is still able to bind and activate EGFR (4). Binding of soluble or membrane bound TGF-alpha to EGFR leads to receptor dimerization, tyrosine autophosphorylation, and activation of downstream signaling components. TGF-alpha and related peptides play an important role in the progression of cancer as well as in neuropathological processes (5,6).

To place an order, please [Click HERE](#).