www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## CD4

Reactivity: Human Mouse

Tested applications: WB IHC FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 FC1:20 - 1:50

Calculated MW:55kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human CD4

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CD4mut;

Polyclonal Antibody

Species: Rabbit

Gene ID:920

Isotype:IgG

Purity: Affinity purification

Swiss Prot:P01730

For research use only.

## Background:

This gene encodes a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigenes and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene.

To place an order, please Click HERE.





