www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

MAPK12

Reactivity: Human Mouse

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:1000 IHC 1:50 - 1:100

Calculated MW:42kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human MAPK12

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

io

Synonym:

ERK3; ERK6; P38GAMMA; PRKM12; SAPK-3; SAPK3

Catalog #:A0559

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:6300

Isotype: IgG

Swiss Prot:P53778

Purity: Affinity purification

For research use only.

Background:

Activation of members of the mitogen-activated protein kinase family is a major mechanism for transduction of extracellular signals. Stress-activated protein kinases are one subclass of MAP kinases. The protein encoded by this gene functions as a signal transducer during differentiation of myoblasts to myotubes. [provided by RefSeq, Jul 2008]

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a CATALOG CLICK HERE