

## PRKCI

**Reactivity:** Human

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:** 68kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human PRKCI

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

qr

**Synonym:**

DXS1179E; MGC26534; PKCI; nPKC-iota

**Catalog #:** A0618

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 5584

**Isotype:** IgG

**Swiss Prot:** P41743

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a member of the protein kinase C (PKC) family of serine/threonine protein kinases. The PKC family comprises at least eight members, which are differentially expressed and are involved in a wide variety of cellular processes. This protein kinase is calcium-independent and phospholipid-dependent. It is not activated by phorbol esters or diacylglycerol. This kinase can be recruited to vesicle tubular clusters (VTCs) by direct interaction with the small GTPase RAB2, where this kinase phosphorylates glyceraldehyde-3-phosphate dehydrogenase (GAPD/GAPDH) and plays a role in microtubule dynamics in the early secretory pathway. This kinase is found to be necessary for BCL-ABL-mediated resistance to drug-induced apoptosis and therefore protects leukemia cells against drug-induced apoptosis. There is a single exon pseudogene mapped on chromosome X. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**