

## TCL1A

**Reactivity:**Human Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**13kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human TCL1A

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

TCL1A;TCL1 ;

**Catalog #:**A0629

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**8115

**Isotype:**IgG

**Swiss Prot:**P56279

**Purity:**Affinity purification

For research use only.

**Background:**

TCL1 (T cell leukemia 1), MTCP1 and TCL1b belong to the TCL1 proto-oncogene family, and their products are involved in Akt activation during embryonic development, T cell leukemias, prolymphocytic leukemias and B cell lymphomas (1-3). The Akt association domain of TCL1 binds with the PH domain of Akt. The formation of an oligomeric TCL-Akt complex is required for TCL1 coactivator function and results in phosphorylation and activation of Akt . Furthermore, functional analysis indicates that the interaction between TCL1 and Akt promotes translocation of Akt to the nucleus (4-6). These findings are supported by the crystal structure of TCL1, which suggests that TCL1 may participate in molecular transport (7).

**To place an order, please [Click HERE](#).**