www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

MMP7

Reactivity: Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:29kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human MMP7

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

MMP7;MMP-7;MPSL1;PUMP-1 ;

Catalog #:A0695

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:4316

Isotype: IgG

Swiss Prot:P09237

Purity: Affinity purification

For research use only.

Background:

The matrix metalloproteinases (MMPs) are a family of proteases that target many extracellular proteins including other proteases, growth factors, cell surface receptors and adhesion molecules (1). Among the family members, MMP-2, MMP-3, MMP-7 and MMP-9 have been characterized as important factors for normal tissue remodeling during embryonic development, wound healing, tumor invasion, angiogenesis, carcinogenesis and apoptosis (2-4). MMP activity correlates with cancer development (2). One mechanism of MMP regulation is transcriptional (5). Once synthesized, MMP exists as a latent proenzyme. Maximum MMP activity requires proteolytic cleavage to generate active MMPs by releasing the inhibitory propeptide domain from the full length protein (5).

To place an order, please Click HERE.

© 2015 NeoScientific