

ATF2

Reactivity: Human Mouse

Tested applications: WB

Recommended Dilution: WB 1:200 - 1:500

Calculated MW: 23kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human ATF2

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fk

Synonym:

ATF2; CRE-BP1; CREB2; HB16; MGC111558; TREB7

Catalog #: A0756

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 1386

Isotype: IgG

Swiss Prot: P15336

Purity: Affinity purification

For research use only.

Background:

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins (1). ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways (2-4). Various forms of cellular stress, including genotoxic agents, inflammatory cytokines, and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71 (2-4). Both SAPK and p38 MAPK have been shown to phosphorylate ATF-2 at these sites in vitro and in cells transfected with ATF-2. Mutations of these sites result in the loss of stress-induced transcription by ATF-2 (2-4). In addition, mutations at these sites reduce the ability of E1A and Rb to stimulate gene expression via ATF-2 (2).

To place an order, please [Click HERE](#).