

EIF2S1

Reactivity:Human Mouse

Tested applications:WB IHC IF IP FC

Recommended Dilution:WB 1:500 - 1:1000 IHC 1:50 - 1:100 IF 1:20 - 1:50 IP 1:20 - 1:50

FC 1:20 - 1:50

Calculated MW:36kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant Protein of human EIF2S1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

EIF2S1;EIF-2;EIF-2A;EIF-2alpha;EIF2;EIF2A ;

Catalog #:A0764

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:1965

Isotype:IgG

Swiss Prot:P05198

Purity:Affinity purification

For research use only.

Background:

Phosphorylation of the eukaryotic initiation factor 2 (eIF2) subunit is a well-documented mechanism to downregulate protein synthesis under a variety of stress conditions. eIF2 binds GTP and Met-tRNAⁱ and transfers Met-tRNA to the 40S subunit to form the 43S preinitiation complex (1,2). eIF2 promotes a new round of translation initiation by exchanging GDP for GTP, a reaction catalyzed by eIF2B (1,2). Kinases that are activated by viral infection (PKR), endoplasmic reticulum stress (PERK/PEK), amino acid deprivation (GCN2), or heme deficiency (HRI) can phosphorylate the subunit of eIF2 (3,4). This phosphorylation stabilizes the eIF2-GDP-eIF2B complex and inhibits the turnover of eIF2B. Induction of PKR by IFN- and TNF- induces potent phosphorylation of eIF2 at Ser51 (5,6).

To place an order, please [Click HERE](#).