NCOA1

Reactivity: Human

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:1000

Calculated MW:157kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human NCOA1

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

Synonym:

F-SRC-1; KAT13A; MGC129719; MGC129720; NCoA-1; RIP160; SRC-1; SRC1; bHLHe42;

bHLHe74

Background:

The protein encoded by this gene acts as a transcriptional coactivator for steroid and nuclear hormone receptors. It is a member of the p160/steroid receptor coactivator (SRC) family and like other family members has histone acetyltransferase activity and contains a nuclear localization signal, as well as bHLH and PAS domains. The product of this gene binds nuclear receptors directly and stimulates the transcriptional activities in a hormone-dependent fashion. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:8648

Isotype:IgG

Swiss Prot:Q15788

Purity: Affinity purification

For research use only.





