www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## **AICDA**

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:24kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human AICDA

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

Synonym:

AID; ARP2; CDA2; HIGM2

Background:

This gene encodes a RNA-editing deaminase that is a member of the cytidine deaminase family. The protein is involved in somatic hypermutation, gene conversion, and class-switch recombination of immunoglobulin genes. Defects in this gene are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2). [provided by RefSeq, Feb 2009]

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:57379 Isotype:IgG

Swiss Prot:Q9GZX7

Purity: Affinity purification

For research use only.



