www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## FCRL2

Reactivity: Human

Tested applications:WB

Recommended Dilution: WB 1:1000 - 1:2000

Calculated MW:56kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human FCRL2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

FCRH2; IFGP4; IRTA4; SPAP1; CD307b; SPAP1A; SPAP1B; SPAP1C;

Background:

This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein has four extracellular C2-type immunoglobulin domains, a transmembrane domain and a cytoplasmic domain that contains one immunoreceptor-tyrosine activation motif and two immunoreceptor-tyrosine inhibitory motifs. This protein may be a prognostic marker for chronic lymphocytic leukemia. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

To place an order, please Click HERE.



Catalog #:A10324

**Antibody Type:** 

Polyclonal Antibody

Species:Rabbit

Gene ID:79368

Isotype:IgG

Swiss Prot:Q96LA5 Purity: Affinity purification

For research use only.





