

STC2

Reactivity:Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:33kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human STC2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

STC-2; STCRP

Catalog #:A10397

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8614

Isotype:IgG

Swiss Prot:O76061

Purity:Affinity purification

For research use only.

Background:

This gene encodes a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The encoded protein has 10 of its 15 cysteine residues conserved among stanniocalcin family members and is phosphorylated by casein kinase 2 exclusively on its serine residues. Its C-terminus contains a cluster of histidine residues which may interact with metal ions. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Constitutive overexpression of human stanniocalcin 2 in mice resulted in pre- and postnatal growth restriction, reduced bone and skeletal muscle growth, and organomegaly. Expression of this gene is induced by estrogen and altered in some breast cancers.

To place an order, please [Click HERE](#).