

SNAI2

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW: 30kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human SNAI2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

k

Synonym:

SNAI2;MGC10182;SLUG;SLUGH1;WS2D ;

Catalog #: A1057

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 6591

Isotype: IgG

Swiss Prot: O43623

Purity: Affinity purification

For research use only.

Background:

Slug (SNAI2) is a widely expressed transcriptional repressor and member of the Snail family of zinc finger transcription factors (1). Similar to the related Snail protein, Slug binds to the E-cadherin promoter region to repress transcription during development (2). The binding of Slug to integrin promoter sequences represses integrin expression and results in reduced cell adhesion (3). Down regulation of E-cadherin expression occurs during the epithelial-mesenchymal transition during embryonic development, a process also exploited by invasive cancer cells (4,5). The tumor suppressor protein p53 induces Slug expression in -irradiated cells; Slug protects damaged cells from apoptosis by repressing p53-induced transcription of the proapoptotic Bcl-2 family protein Puma (6). Deletion mutations in the corresponding Slug gene are associated with the pigmentation disorders Waardenburg Syndrome and Piebaldism, while a genetic duplication resulting in Slug overexpression is associated with a collection of congenital heart defects termed tetralogy of Fallot (7).

To place an order, please [Click HERE](#).