

YWHAE

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:100

Calculated MW:29kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human YWHAE

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fjng

Synonym:

YWHAE;14-3-3E;FLJ45465;KCIP-1;MDCR;MDS ;

Catalog #:A1058

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7531

Isotype:IgG

Swiss Prot:P62258

Purity:Affinity purification

For research use only.

Background:

The 14-3-3 family of proteins plays a key regulatory role in signal transduction, checkpoint control, apoptotic and nutrient-sensing pathways (1,2). 14-3-3 proteins are highly conserved and ubiquitously expressed. There are at least seven isoforms, γ , δ , ϵ , ζ , η , θ , and ι that have been identified in mammals. The initially described δ and ϵ isoforms are confirmed to be phosphorylated forms of δ and ϵ , respectively (3). Through their amino-terminal helical region, 14-3-3 proteins form homo- or heterodimers that interact with a wide variety of proteins: transcription factors, metabolic enzymes, cytoskeletal proteins, kinases, phosphatases, and other signaling molecules (3,4). The interaction of 14-3-3 proteins with their targets is primarily through a phospho-Ser/Thr motif. However, binding to divergent phospho-Ser/Thr motifs, as well as phosphorylation-independent interactions, has been observed (4). 14-3-3 binding masks specific sequences of the target protein and therefore modulates target protein localization, phosphorylation state, stability, and molecular interactions (1-4). 14-3-3 proteins may also induce target protein conformational changes that modify target protein function (4,5). Distinct temporal and spatial expression patterns of 14-3-3 isoforms have been observed in development and in acute response to extracellular signals and drugs, suggesting that 14-3-3 isoforms may perform different functions despite their sequence similarities (4). Several studies suggest that 14-3-3 isoforms are differentially regulated in cancer and neurological syndromes (2,3).

To place an order, please [Click HERE](#).