

## ACTR3

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**47kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human ACTR3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ACTR3;ARP3 ;

**Catalog #:**A1064

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**10096

**Isotype:**IgG

**Swiss Prot:**P61158

**Purity:**Affinity purification

For research use only.

**Background:**

Actin nucleation, the formation of new actin filaments from existing filaments, affects actin filament structure during cell motility, division, and intracellular trafficking. An important actin nucleation protein complex is the highly conserved ARP2/3 complex, consisting of ARP2, ARP3, and ARPC1-5. The ARP2/3 complex promotes branching of an existing actin filament and formation of a daughter filament following activation by nucleation-promoting factors, such as WASP/WAVE or cortactin (1). The formation of podosomes, small cellular projections that degrade the extracellular matrix, is enhanced by ARP2/3 complex action. ARP2/3 competes with caldesmon, an actin binding protein shown to negatively affect podosome formation (2). Along with N-WASP, the ARP2/3 complex regulates nuclear actin filament nucleation and controls actin polymerization during transcription (3).

*To place an order, please [Click HERE](#).*