

## UBE2K

**Reactivity:** Human Mouse Rat

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:2000

**Calculated MW:** 22kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human UBE2K

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

fl

**Synonym:**

UBE2K;DKFZp564C1216;DKFZp686J24237;E2-25K;HIP2;HYPG;LIG ;

**Catalog #:** A1086

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 3093

**Isotype:** IgG

**Swiss Prot:** P61086

**Purity:** Affinity purification

For research use only.

**Background:**

Protein ubiquitination requires the concerted action of the E1, E2, and E3 ubiquitin-conjugating enzymes. Ubiquitin is first activated through ATP-dependent formation of a thiol ester with ubiquitin-activating enzyme E1. The activated ubiquitin is then transferred to a thiol group of ubiquitin-carrier enzyme E2. The final step is the transfer of ubiquitin from E2 to an -amino group of the target protein lysine residue, which is mediated by ubiquitin-ligase enzyme E3 (1). E2-25K (Hip2) is a member of the E2 protein family that catalyzes multiubiquitin chain synthesis via Lys48 of ubiquitin (2). E2-25K is reportedly involved in Alzheimer's disease, Huntington's disease and antigen processing through its interaction with amyloid-, huntingtin, and MHC-heavy chain proteins (3-5). Lys14 of E2-25K can be modified by SUMOylation, with this modification resulting in inhibited E2 activity (6).

**To place an order, please [Click HERE](#).**