

## NGFR

---

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC ICC IF IP

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100  
IP 1:20 - 1:50

**Calculated MW:**75 kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human NGFR

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR;

**Catalog #:**A11050

**Antibody Type:**

Monoclonal Antibody

**Species:**Rabbit

**Gene ID:**4804

**Isotype:**IgG

**Swiss Prot:**P08138

**Purity:**Affinity purification

For research use only.

**Background:**

Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

*To place an order, please [Click HERE](#).*