

NOTCH1

Reactivity:Human Mouse

Tested applications:WB IHC ICC IF IP FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100
IP 1:20 - 1:50 FC 1:20 - 1:50

Calculated MW:125 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human NOTCH1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

hN1; AOS5; TAN1; AOVD1;

Catalog #:A11056

Antibody Type:

Monoclonal Antibody

Species:Rabbit

Gene ID:4851

Isotype:IgG

Swiss Prot:P46531

Purity:Affinity purification

For research use only.

Background:

This gene encodes a member of the NOTCH family of proteins. Members of this Type I transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple different domain types. Notch signaling is an evolutionarily conserved intercellular signaling pathway that regulates interactions between physically adjacent cells through binding of Notch family receptors to their cognate ligands. The encoded preproprotein is proteolytically processed in the trans-Golgi network to generate two polypeptide chains that heterodimerize to form the mature cell-surface receptor. This receptor plays a role in the development of numerous cell and tissue types. Mutations in this gene are associated with aortic valve disease, Adams-Oliver syndrome, T-cell acute lymphoblastic leukemia, chronic lymphocytic leukemia, and head and neck squamous cell carcinoma.

To place an order, please [Click HERE](#).