

MAP2K2

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IP

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IP 1:20 - 1:50

Calculated MW: 44 kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human MAP2K2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CFC4; MEK2; MKK2; MAPKK2; PRKMK2;

Catalog #: A11122

Antibody Type:

Monoclonal Antibody

Species: Rabbit

Gene ID: 5605

Isotype: IgG

Swiss Prot: P36507

Purity: Affinity purification

For research use only.

Background:

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

To place an order, please [Click HERE](#).