www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

VCAM1

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IF IP

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100

IP 1:20 - 1:50

Calculated MW:100 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human VCAM1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

CD106; INCAM-100;

Background:

This gene is a member of the Ig superfamily and encodes a cell surface sialoglycoprotein expressed by cytokine-activated endothelium. This type I membrane protein mediates leukocyte-endothelial cell adhesion and signal transduction, and may play a role in the development of artherosclerosis and rheumatoid arthritis. Three alternatively spliced transcripts encoding different isoforms have been described for this gene.

To place an order, please Click HERE.

Antibody Type:

Monoclonal Antibody

Species: Rabbit

Gene ID:7412

Isotype:IgG

Swiss Prot:P19320 Purity: Affinity purification

For research use only.



