

TNNT2

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:36kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant Protein of human TNNT2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

TNNT2 ; Cardiac muscle troponin T; Troponin T; cardiac muscle; troponin T type 2 (cardiac); cTnT; TnTC

Catalog #:A1126

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7139

Isotype:IgG

Swiss Prot:P45379

Purity:Affinity purification

For research use only.

Background:

Troponin, working in conjunction with tropomyosin, functions as a molecular switch, regulating muscle contraction in response to changes in the intracellular Ca²⁺ concentration. Troponin consists of three subunits: the Ca²⁺-binding subunit troponin C (TnC), the tropomyosin-binding subunit troponin T (TnT), and the inhibitory subunit troponin I (TnI) (1). In response to -adrenergic stimulation of the heart, Ser23 and Ser24 of TnI (cardiac) are phosphorylated by PKA and PKC. This phosphorylation stimulates a conformational change of the regulatory domain of TnC, reduces the association between TnI and TnC, and decreases myofilament Ca²⁺ sensitivity by reducing the Ca²⁺ binding affinity of TnC (1-3). The tropomyosin binding subunit of the troponin complex TnT exists as different isoforms in slow skeletal muscle (ssTnT/TNNT1), fast skeletal muscle (fsTnT/TNNT3) and in cardiac muscle (cTnT/TNNT2). Each of these may also contain multiple alternatively spliced variants. Assays for measuring serum concentrations of cTnT, as well as cTnI, have been reported for analyzing cardiac injury.

To place an order, please [Click HERE](#).