

## F3

**Reactivity:**Human Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**33kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human F3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

TF; TFA; CD142; F3;

**Catalog #:**A1378

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**2152

**Isotype:**IgG

**Swiss Prot:**P13726

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010]

**To place an order, please [Click HERE](#).**