

## PSMC5

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**46kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human PSMC5

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

S8; p45; SUG1; SUG-1; TBP10; TRIP1; p45/SUG;PSMC5;

**Background:**

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits.

Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. In addition to participation in proteasome functions, this subunit may participate in transcriptional regulation since it has been shown to interact with the thyroid hormone receptor and retinoid X receptor-alpha. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2010]

**To place an order, please [Click HERE](#).**

**Catalog #:**A1538

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5705

**Isotype:**IgG

**Swiss Prot:**P62195

**Purity:**Affinity purification

For research use only.