

B2M

Reactivity:Human Mouse

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:14kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human B2M

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

beta-2-microglobulin; B2M;

Background:

2-microglobulin (B2M) is a principal component of the Major Histocompatibility Complex (MHC) class I molecule, a ternary membrane protein complex that displays fragments derived from proteolyzed cytosolic proteins on the surface of cells for recognition by the surveillance immune system (1,2). As an integral component of the MHC class I complex, 2-microglobulin plays a critically important role in immune system function (3). It has important relevance to cancer biology research; for example, research studies have shown that nearly one-third of diffuse large B cell lymphomas contain mutations that inactivate 2-microglobulin gene function, thereby allowing tumor cells to escape immune detection (4). In addition, 2-microglobulin has been identified as an amyloid preprotein with collagen-binding affinity (5); its accumulation in osteoarthritic lesions of long-term dialysis patients is reportedly a contributing factor to the condition known as amyloid osteoarthropathy (6).

To place an order, please [Click HERE](#).

Catalog #:A1562

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:567

Isotype:IgG

Swiss Prot:P61769

Purity:Affinity purification

For research use only.